



The National Comprehensive Cancer Network (NCCN [v 3.2019](#) and [v1.2020](#)) guidelines recommend referral to a cancer genetics professional for an individual with one or more of the following:

AFFECTED WITH OVARIAN CANCER

- Epithelial ovarian cancer (including Fallopian tube cancer or peritoneal cancer) at any age

FAMILY HISTORY OF OVARIAN CANCER

- First- or second-degree relative with epithelial ovarian cancer (including Fallopian tube cancer or peritoneal cancer) at any age

**Referral is also clinically indicated for individuals meeting criteria but with previous limited testing (eg, single gene and/or absent deletion duplication analysis) interested in pursuing multi-gene testing*

CONDITIONS TO CONSIDER:

- [BRCA-related cancer susceptibility](#)
- BRIP1 related cancer susceptibility
- DICER1 related cancer susceptibility (Non-epithelial ovarian cancers)
- [EPCAM-related cancer susceptibility](#)
- [MLH1-related cancer susceptibility](#)
- [MSH2-related cancer susceptibility](#)
- [MSH6-related cancer susceptibility](#)
- [PMS2-related cancer susceptibility](#)
- RAD51C-related cancer susceptibility
- RAD51D-related cancer susceptibility
- [STK11-related cancer susceptibility](#) (Non-epithelial ovarian cancers)